
Purpose: The objective of this report is to provide timely information regarding the status of opioids including novel opioids and fentanyl analog prevalence, positivity, and discovery within the United States.

Project Overview: There are many indicators from emergency room admissions, poison center calls, drug treatment admissions, drug possession and trafficking arrests, crime laboratory statistics, and medical examiner data, that the illicit use and abuse of both prescription opioids such as morphine, fentanyl and oxycodone, and traditionally abused opioids, most notably heroin are increasing. In addition, many novel opioid agonists have been identified over the last three years that are now contributing to the opioid death statistics at an increasing but poorly documented rate. In collaboration with NMS Labs, this project sought to use de-identified raw electronic high resolution mass spectrometry data from postmortem and driving under the influence of drugs (DUID) case to develop trend reports and perform datamining and repeated interrogations of the raw data, against a continually updated database of common opioids and emerging opioid drugs. Data processed against a library containing 170 different opioids and/or fentanyl analogs.

Opioid Positivity

Between January and March 2019, there were a total of 20,014 cases positive for one or more opioids that are part of routine testing. Fentanyl accounted for the highest positivity with 4,353 positive cases followed by heroin with 1,621 and morphine with 1,666.

OPIOID POSITIVITY JANUARY - MARCH 2019



^{*}Heroin positivity determined by detecting 6-MAM in any fluid and morphine in the blood.

Fentanyl



Data has not been normalized to account for testing volume. Geographical distribution may be limited by location of agencies submitting samples to NMS Labs and not necessarily representative of all jurisdictions.

Fentanyl Demographics (n= 4,353)

Sex			
Male	2,681		
Female	1,050		
Unknown	622		
Age			
Average	40 (±13)		
Median	43		
Range	0-96		

82% of cases were death investigation; 14% were law enforcement


Heroin

Heroin Demographics (n=1,621)

Sex		
Male	1,116	
Female	343	
Unknown	162	
Age		
Average	40 (±12)	
Median	43	
Range	18-82	

93% of cases were death investigation; 4% were law enforcement

Heat Map of Heroin Positive Cases January - March 2019



Data has not been normalized to account for testing volume. Geographical distribution may be limited by location of agencies submitting samples to NMS Labs and not necessarily representative of all jurisdictions.








Acknowledgements: This report was prepared by Amanda L.A. Mohr, MSFS, D-ABFT-FT; Melissa F. Fogarty MSFS, D-ABFT-FT; Judith Rodriguez Salas MSFS; and Barry K. Logan, PhD, F-ABFT at the Center for Forensic Science Research and Education at the Fredric Rieders Family Foundation. This project is supported by Award Number 2017-DN-BX-0169, awarded by the National Institute of Justice, Office of Justice Programs, U.S. Department of Justice. The opinions, findings, and conclusions or recommendations expressed in this publication are those of the authors and do not necessarily reflect those of the Department of Justice.

Novel Opioid Positivity


Between January and March 2019, there were a total of 221 cases positive for one or more novel opioid. Para-fluoroisobutyrylfentanyl (p-FIBF) accounted for the highest positivity with 74 cases followed by valerylfentanyl with 44 cases and carfentanil with 23 cases.

NOVEL OPIOID POSITIVITY JANUARY - MARCH 2019



Novel Opioid o/m/p-Fluorofuranylfentanyl Identified in January 2019


Retrospectively identified two cases from December 2018



Case Receipt	Demographics		ohics	Analytes Identified in Addition to	
Date	Age	Sex	State	o/m/p-Fluorofuranylfentanyl	
December 2018	N/A	F	FL	Naloxone, Methamphetamine, Alprazolam, Cocaine, Benzoylecgonine, Hydromorphone, Nortriptyline, THC, Citalopram	
December 2018	62	F	PA	Naloxone, Oxazepam, Temazepam, Lorazepam, Morphine, 6-MAM, Fentanyl	
January 2019	34	M	NJ	Ethanol, Naloxone, p-FIBF, Alprazolam	
January 2019	27	M	NJ	Naloxone, THC	
March 2019	39	M	NJ	THC, Amphetamine, Methamphetamine, p-FIBF	

Novel Opioids

Heat Map of Novel Opioid Cases January - March 2019



Data has not been normalized to account for testing volume. Geographical distribution may be limited by location of agencies submitting samples to NMS Labs and not necessarily representative of all jurisdictions.

Demographic Information for Top Three Novel Opioids

p-FIBF Demographics (n=74)

Sex			
Male	19		
Female	50		
Unknown	5		
Age			
Average	37 (±10)		
Median	40		
Range	17-56		

Valerylfentanyl Demographics (n=44)

Sex				
Male	27			
Female	12			
Unknown	5			
Age				
Average	41 (±13)			
Median	40			
Range	21-68			

Carfentanil
Demographics (n= 23)

Sex			
Male	14		
Female	7		
Unknown	2		
Age			
Average	36 (±11)		
Median	40		
Range	19-54		

87% of cases were death investigation cases; 8% were law enforcement; 3% were hospital





